

DATE: \_\_\_\_\_

PATIENT NAME: \_\_\_\_\_

AGE: \_\_\_\_\_ SEX: \_\_\_\_\_

BREED: \_\_\_\_\_ WEIGHT: \_\_\_\_\_


**Right Eye**

**Prescription**

**Left Eye**

**Treatment Time**

**Treatment Time**



1. Eye wash
  - Rinse both eyes of mucus  $\geq 2$  times per day.
  - Use a water-soaked cotton ball to wipe debris from eyelids.
  - Use separate cotton for each eye and discard.

**Wait 1 to 2 minutes**

2. Betadine solution (1:5 dilution)
  - Mix 5 mL of water with 0.1 mL of betadine.
  - Using a saturated cotton ball, instill a few drops in the right eye.
  - Cleanse lids 2 times per day.
  - Mix fresh solution for each administration. Solution can stain.

**Wait 5 to 10 minutes**

3. Tobramycin 0.3% ophthalmologic solution (antibiotic)
  - Instill 1 drop in right eye 4 times per day.

**Wait 5 to 10 minutes**

4. Autologous serum
  - Keep refrigerated.
  - Instill 1 drop in the right eye 4 times per day.
  - Discard after 5 days.
  - Refill as needed.

**Wait 5 to 10 minutes**

5. Ofloxacin 0.3% ophthalmologic solution (antibiotic)
  - Instill 1 drop in right eye 4 times per day.

6. Doxycycline (100-mg chews)
  - Give chew orally 2 times per day with plenty of water.
  - Do not give with cheese or calcium-containing products.

Follow chart carefully. Wait between medication administrations as indicated. Keep Elizabethan collar on at all times. Offer food and water, and monitor eating. Use a soft, water-moistened paper towel to wipe the inside of the Elizabethan collar as needed to keep clean of food and drool. If you think the eye(s) is getting worse or observe any change (eg, more painful, more discharge, squinting, restless, loss of appetite), contact your veterinarian immediately.

Please bring all medications and checklists to your next appointment. NEXT APPOINTMENT: \_\_\_\_\_